WHAT'S AHEAD

- **1.1** The Study of Chemistry
- **1.2** Classifications of Matter
- **1.3** Properties of Matter
- **1.4** The Nature of Energy
- **1.5 Units of Measurement**
- **1.6 Uncertainty in Measurement**
- **1.7** Dimensional Analysis

INTRODUCTION: MATTER, ENERGY, AND MEASUREMENT

1.1 | The Study of Chemistry



The title of this book—*Chemistry: The Central Science*—reflects the fact that much of what goes on in the world around us involves chemistry. Everyday chemical processes include the changes that produce the brilliant colors of flowers, the ways our bodies process the food we eat, and the electrical energy that powers our cell phones.

This first chapter provides an overview of what chemistry is about and what chemists do. The "What's Ahead" list gives an overview of the chapter organization and of some of the ideas we will consider.

Chemistry is the study of matter, its properties, and the changes that matter undergoes. As you progress in your study, you will come to see how chemical principles operate in all aspects of our lives, from everyday activities like food preparation to more complex processes such as those that operate in the environment. We will also learn how the properties of substances can be tailored for specific applications by controlling their composition and structure. For example, the synthetic pigments chemists developed in the nineteenth century were used extensively by impressionist artists such as van Gogh.

Chemistry is at the heart of many changes we see in the world around us, and it accounts for the myriad different properties we see in matter. To understand how these changes and properties arise, we need to look far beneath the surfaces of our everyday observations.

By the end of this section, you should be able to

• Appreciate the scope of chemistry

The Atomic and Molecular Perspective of Chemistry

Chemistry is the study of the properties and behavior of matter. **Matter** is the physical material of the universe; it is anything that has mass and occupies space. A **property** is any characteristic that allows us to recognize a particular type of matter and to distinguish it from other types. This book, your body, the air you are breathing, and the clothes you are wearing are all samples of matter. We observe a tremendous variety of matter in our world, but countless experiments have shown that all matter is comprised of combinations of only about 100 substances called **elements**. One of our major goals will be to relate the properties of matter to its composition, that is, to the particular elements it contains.

Chemistry also provides a background for understanding the properties of matter in terms of **atoms**, the almost infinitesimally small building blocks of matter. Each element is composed of a unique kind of atom. We will see that the properties of matter relate to both the kinds of atoms the matter contains (*composition*) and the arrangements of these atoms (*structure*).

In **molecules**, two or more atoms are joined in specific shapes. Throughout this text you will see molecules represented using colored spheres to show how the atoms are connected (Figure 1.1). The color provides a convenient way to distinguish between atoms of different elements. For example, notice that the molecules of ethanol and ethylene glycol in Figure 1.1 have different compositions and structures. Ethanol contains one oxygen atom, depicted by one red sphere. In contrast, ethylene glycol contains two oxygen atoms.

Even apparently minor differences in the composition or structure of molecules can cause profound differences in properties. For example, let's compare ethanol and ethylene glycol, which appear in Figure 1.1 to be quite similar. Ethanol is the alcohol in beverages such as beer and wine, whereas ethylene glycol is a viscous liquid used as automobile antifreeze. The properties of these two substances differ in many ways, as do their biological activities. Ethanol is consumed throughout the world, but you should *never* consume ethylene glycol because it is highly toxic. One of the challenges chemists undertake is to alter the composition or structure of molecules in a controlled way, creating new substances with different properties. For example, the common drug aspirin, shown in Figure 1.1, was first synthesized in 1897 in a successful attempt to



improve on a natural product extracted from willow bark that had long been used to alleviate pain.

Every change in the observable world—from boiling water to the changes that occur as our bodies combat invading viruses—has its basis in the world of atoms and molecules. Thus, as we proceed with our study of chemistry, we will find ourselves thinking in three realms: the *macroscopic* realm of ordinary-sized objects (*macro* = large), the *microscopic* realm of atoms and molecules, and the symbolic realm of how we represent these particles. We make our observations in the macroscopic world, but to understand that world, we must visualize how atoms and molecules behave at the microscopic level. Chemistry is the science that seeks to understand the properties and behavior of matter by studying the properties and behavior of atoms and molecules.

Why Study Chemistry?

Chemistry lies near the heart of many matters of public concern, such as improvement of health care, conservation of natural resources, protection of the environment, and the supply of energy needed to keep society running. Using chemistry, we have discovered and continually improved upon pharmaceuticals, fertilizers and pesticides, plastics, solar panels, light-emitting diodes (LEDs), and building materials. We have also discovered that some chemicals are harmful to our health or the environment. This means that we must be sure that the materials with which we come into contact are safe. As a citizen and consumer, it is in your best interest to understand the effects, both positive and negative, that chemicals can have, in order to arrive at a balanced outlook regarding their uses.

You may be studying chemistry because it is an essential part of your curriculum. Your major might be chemistry, or it could be biology, engineering, pharmacy, agriculture, geology, or some other field. Chemistry is central to a fundamental understanding of governing principles in many science-related fields. For example, our interactions with the material world raise basic questions about the materials around us. **Figure 1.2** illustrates how chemistry is central to several different realms of modern life.



▲ Figure 1.2 Chemistry is central to our understanding of the world around us.

CHEMISTRY PUT TO WORK Chemistry and the Chemical Industry

Chemistry is all around us. For example, sugar is a chemical extracted from natural sources, refined by industry and sold across the world in a highly purified form. However, few realize the size and importance of the chemical industry. The chemical industry in the United States is estimated to be an \$800 billion enterprise that employs over 800,000 people and accounts for 14% of all U.S. exports.

Who are chemists, and what do they do? People who have degrees in chemistry hold a variety of positions in industry, government, and academia. Those in industry work as laboratory chemists, developing new products (research and development); analyzing materials (quality control); or assisting customers in using products (sales and service). Those with more experience or training may work as managers or company directors. Chemists are important members of the scientific workforce in government (the National Institutes of Health, Department of Energy, and Environmental Protection Agency all employ chemists) and at universities. A chemistry degree is also good preparation for careers in teaching, medicine, biomedical research, information science, environmental work, technical sales, government regulatory agencies, and patent law.

Fundamentally, chemists do three things: They (1) make new types of matter: materials, substances, or combinations of substances with desired properties; (2) measure the properties of matter; and (3) develop models that explain and/or predict the properties of matter. One chemist, for example, may work in the laboratory to discover new drugs. Another may concentrate on the development of new instrumentation to measure properties of matter at the atomic level. Other chemists may use existing materials and methods to understand how pollutants are transported in the environment or how drugs are processed in the body. Yet another chemist will develop theory, write computer code, and run computer simulations to understand how molecules move and react. The collective chemical enterprise is a rich mix of all of these activities.



▲ Figure 1.3 Common chemicals used for household cleaning.

Self-Assessment Exercise

- **1.1** Which of the following groups of substances involve the use of chemicals? Indicate all that apply.
 - (a) Paints, printer toner, food coloring
 - (b) Computer displays, LED lights, barcode readers

- (c) Antiseptic cream, pain killers, energy drinks
- (d) A light-weight bicycle frame, food packaging, a car exhaust catalytic converter
- (e) Soap, shampoo, washing powder

ment to IIA 1.1

🔬 Answers to Self-Assessment Exercises



The study of any science starts with classification. This brings order to what we study much as the sorting of books in a library enables you to find the book you want. Let's begin our study of chemistry by examining two fundamental ways in which matter is classified. Matter is typically characterized by (1) its physical state (gas, liquid, or solid) and (2) its composition (whether it is an element, a *compound*, or a *mixture*). By the end of this section, you should be able to

· Understand the basic classification of matter

States of Matter

A sample of matter can be a gas, a liquid, or a solid. These three forms, called the **states of matter**, differ in some of their observable properties.

- A **gas** (also known as vapor) has no fixed volume or shape; rather, it uniformly fills its container. A gas can be compressed to occupy a smaller volume, or it can expand to occupy a larger one.
- A **liquid** has a distinct volume independent of its container, assumes the shape of the portion of the container it occupies, and is not compressible to any appreciable extent.
- A **solid** has both a definite shape and a definite volume and is not compressible to any appreciable extent.

The properties of the states of matter can be understood on the molecular level (Figure 1.4). In a gas the molecules are far apart and moving at high speeds, colliding repeatedly with one another and with the walls of the container. Compressing a gas decreases the amount of space between molecules and increases the frequency of collisions between molecules

1.2 Classifications of Matter

In which form of water are the water molecules

but does not alter the size or shape of the molecules. In a liquid, the molecules are packed closely together but still move rapidly. The rapid movement allows the molecules to slide over one another; thus, a liquid pours easily. In a solid the molecules are held tightly together, usually in definite arrangements in which the molecules can wiggle only slightly in their otherwise fixed positions. Thus, the distances between molecules are similar in the liquid and solid states, but while the molecules are for the most part locked in place in a solid, they retain considerable freedom of motion in a liquid. Changes in temperature and/or pressure can lead to conversion from one state of matter to another, illustrated by such familiar processes as ice melting or water vapor condensing.

Pure Substances

Most forms of matter we encounter—the air we breathe (a gas), the fuel we burn in our cars (a liquid), and the road they run on (a solid)—are not chemically pure. We can, however, separate these forms of matter into pure substances. A **pure substance** (usually referred to simply as a *substance*) is matter that has distinct properties and a composition that does not vary from sample to sample. Water and table salt (so-dium chloride) are examples of pure substances.

All substances are either elements or compounds.

• **Elements** are substances that cannot be decomposed into simpler substances. On the molecular level, each element is composed of only one kind of atom [Figure 1.5(a and b)].



Figure 1.5(**d**) shows a mixture of substances. **Mixtures** are combinations of two or more substances in which each substance retains its chemical identity.





Go Figure

▲ Figure 1.4 The three physical states of water—water vapor, liquid water, and ice. We see the liquid and solid states but cannot see the gas (vapor) state. The red arrows show that the three states of matter interconvert.

Go Figure

If the lower pie chart was drawn as the percentage in terms of number of atoms rather than the percentage in terms of mass, would the hydrogen slice of the pie get larger or smaller?



▲ Figure 1.6 Relative abundances of elements.* Elements in percent by mass in Earth's crust (including oceans and atmosphere) and the human body.

Elements

Currently, 118 elements are known, though they vary widely in abundance. Hydrogen constitutes about 74% of the mass in the Milky Way galaxy, and helium constitutes 24%. Closer to home, only five elements—oxygen, silicon, aluminum, iron, and calcium—account for over 90% of Earth's crust (including oceans and atmosphere), and only three—oxygen, carbon, and hydrogen account for over 90% of the mass of the human body (Figure 1.6).

Table 1.1 lists some common elements, along with the chemical *symbols* used to denote them. The symbol for each element consists of one or two letters, with the first letter capitalized. These symbols are derived mostly from the English names of the elements, but sometimes they are derived from a foreign name instead (last column in Table 1.1). You will need to know these symbols and learn others as we encounter them in the text.

All of the known elements and their symbols are listed on the front inside cover of this text in a table known as the *periodic table*. In the periodic table, the elements are arranged in columns so that closely related elements are grouped together. We describe the periodic table in more detail in Section 2.5 and consider the periodically repeating properties of the elements in Chapter 7.

Compounds

Most elements can interact with other elements to form compounds. For example, when hydrogen gas burns in oxygen gas, the elements hydrogen and oxygen combine to form the compound water. Conversely, water can be decomposed into its elements by passing an electrical current through it (Figure 1.7).

Decomposing pure water into its constituent elements shows that it contains 11% hydrogen and 89% oxygen by mass, regardless of its source. This ratio is constant because every water molecule has the same number of hydrogen and oxygen atoms. While the mass percentages make it seem that water is mostly oxygen, there are actually two hydrogen atoms and only one oxygen atom per molecule. The explanation for this apparent discrepancy comes from the fact that hydrogen atoms are much lighter than oxygen atoms. The mass composition, a macroscopic determination, corresponds to

TABLE 1.1 Some Common Elements and Their Symbols

Carbon	С	Aluminum	Al	Copper	Cu (from <i>cuprum</i>)	
Fluorine	F	Bromine	Br	Iron	Fe (from <i>ferrum</i>)	
Hydrogen	Н	Calcium	Ca	Lead	Pb (from <i>plumbum</i>)	
Iodine	Ι	Chlorine	Cl	Mercury	Hg (from <i>hydrargyrum</i>)	
Nitrogen	Ν	Helium	He	Potassium	K (from kalium)	
Oxygen	0	Lithium	Li	Silver	Ag (from <i>argentum</i>)	
Phosphorus	Р	Magnesium	Mg	Sodium	Na (from natrium)	
Sulfur	S	Silicon	Si	Tin	Sn (from <i>stannum</i>)	

^{*}U.S. Geological Survey Circular 285, U.S Department of the Interior.



As seen in **Table 1.2**, the properties of water bear no resemblance to the properties of its component elements. Hydrogen, oxygen, and water are each a unique substance, a consequence of the uniqueness of their respective molecules.

Able 1.2 Companison of water, Aydrogen, and Oxygen			
	Water	Hydrogen	Oxygen
State ^a	Liquid	Gas	Gas
Normal boiling point	100 °C	−253 °C	−183 °C
Density ^a	$1000 \ kg/m^3$	$0.084 \ kg/m^3$	$1.33 \ kg/m^3$
Flammable	No	Yes	No

ABLE 1.2	Comparison	of Water,	Hydrogen,	and Oxygen	

^a At room temperature and atmospheric pressure.

The observation that the elemental composition of a compound is always the same is known as the law of constant composition (or the law of definite proportions). French chemist Joseph Louis Proust (1754–1826) first stated the law in about 1800. Although this law has been known for 200 years, the belief persists among some people that a fundamental difference exists between compounds prepared in the laboratory and the corresponding compounds found in nature. This simply is not true. Regardless of its source—nature or a laboratory—a pure compound has the same composition and properties under the same conditions. Both chemists and nature must use the same elements and operate under the same natural laws. When two materials differ in composition or properties, either they are composed of different compounds or they differ in purity.

Mixtures

Most of the matter we encounter consists of mixtures of different substances. Each substance in a mixture retains its chemical identity and properties. In contrast to a pure substance, which by definition has a fixed composition, the composition of a mixture can vary. A cup of sweetened coffee, for example, can contain either a little sugar or a lot. The substances making up a mixture are called *components* of the mixture.

Some mixtures do not have the same composition, properties, and appearance throughout. Rocks and wood, for example, vary in texture and appearance in any typical sample. Such mixtures are *heterogeneous* [Figure 1.8(a)]. Mixtures that are uniform throughout are homogeneous. Air is a homogeneous mixture of nitrogen, oxygen, and smaller amounts of other gases. The nitrogen in air has all the properties of pure nitrogen because both the pure substance and the mixture contain the same nitrogen molecules. Salt, sugar, and many other substances dissolve in water to form **solutions**, which are homogeneous mixtures [Figure 1.8(b)]. Although the term *solution* conjures an image of a liquid, the term can also apply to solids, liquids, or gases.

Figure 1.9 summarizes the classification of matter into elements, compounds, and mixtures.

Figure 1.8 Mixtures. (a) Many common materials, including rocks, are heterogeneous mixtures. This photograph of granite shows a heterogeneous mixture of silicon dioxide and other metal oxides. (b) Homogeneous mixtures are called solutions. Many substances, including the blue solid shown here [copper(II) sulfate pentahydrate], dissolve in water to form solutions.



(a)





Distinguishing among Elements, Compounds, and Mixtures

"White gold" contains gold and a "white" metal, such as palladium. Two samples of white gold differ in the relative amounts of gold and palladium they contain. Both samples are uniform in composition throughout. Use Figure 1.9 to classify white gold.

SOLUTION

Because the material is uniform throughout, it is homogeneous. Because its composition differs for the two samples, it cannot be a compound. Instead, it must be a homogeneous mixture.

Practice Exercise

Aspirin is composed of 60.0% carbon, 4.5% hydrogen, and 35.5% oxygen by mass, regardless of its source. Use Figure 1.9 to classify aspirin.

Self-Assessment Exercise

- **1.2** Carbonated water, 'Soda Water', is water in which carbon dioxide has been dissolved under pressure. Classify soda water.
- (a) A homogeneous solution of elements
- (b) A homogeneous solution of compounds
- (c) A heterogeneous mixture of elements
- (d) A heterogeneous mixture of compounds

Exercises

- 1.3 Classify each of the following as a pure substance or a mixture. If a mixture, indicate whether it is homogeneous or heterogeneous: (a) air, (b) chocolate with almond, (c) aluminium, (d) iodine tincture.
- 1.4 Give the chemical symbol or name for the following elements, as appropriate: (a) helium, (b) platinum, (c) cobalt, (d) tin, (e) silver, (f) Sb, (g) Pb, (h) Br, (i) V, (j) Hg.
- **1.5** A solid white substance A is heated strongly in the absence of air. It decomposes to form a new white substance B and a gas C. The gas has exactly the same properties as the product obtained when carbon is burned in an excess of oxygen. Based on these observations, can we determine whether solids A and B and gas C are elements or compounds?

(d) **S.**r

1.3 Properties of Matter



We are familiar with many of the properties of materials around us—the hardness of steel, the flexibility of rubber, or the glow of a wood fire. If we delve a little deeper, we can classify these properties as one of two types. By the end of this section, you should be able to

· Distinguish between physical and chemical changes

Every substance has unique properties. For example, the properties listed in Table 1.2 allow us to distinguish hydrogen, oxygen, and water from one another. The properties of matter can be categorized as physical or chemical. **Physical properties** can be observed without changing the identity and composition of the substance. These properties include color, odor, density, melting point, boiling point, and hardness. **Chemical properties** describe the way a substance may change, or *react*, to form other substances. A common chemical property is flammability, the ability of a substance to burn in the presence of oxygen.

Some properties, such as temperature and melting point, are *intensive properties*. **Intensive properties** do not depend on the amount of sample being examined and are particularly useful in chemistry because many intensive properties can be used to *identify* substances. **Extensive properties** depend on the amount of sample, with two examples being mass and volume. Extensive properties relate to the *amount* of substance present.

Physical and Chemical Changes

The changes substances undergo are either physical or chemical. During a **physical change**, a substance changes its physical appearance but not its composition. (That is, it is the same substance before and after the change.) The evaporation of water is a physical change. When water evaporates, it changes from the liquid state to the gas state, but it is still composed of water molecules, as depicted in Figure 1.4. All **changes of state** (for example, from liquid to gas or from liquid to solid) are physical changes.

In a **chemical change** (also called a **chemical reaction**), a substance is transformed into a chemically different substance. When hydrogen burns in air, for example, it undergoes a chemical change because it combines with oxygen to form water (Figure 1.10).

Chemical changes can be dramatic. In the account given in **Figure 1.11**, Ira Remsen, author of a popular chemistry text published in 1901, describes his first experiences with chemical reactions.

Separation of Mixtures

We can separate a mixture into its components by taking advantage of differences in their properties. For example, a heterogeneous mixture of iron filings and gold filings could be sorted by color into iron and gold. A less tedious approach would be to use a magnet to



▲ Figure 1.10 A chemical reaction.

While reading a textbook of chemistry, I came upon the statement "nitric acid acts upon copper," and I determined to see what this meant. Having located some nitric acid, I had only to learn what the words "act upon" meant. In the interest of knowledge I was even willing to sacrifice one of the few copper cents then in my possession. I put one of them on the table, opened a bottle labeled "nitric acid," poured some of the liquid on the copper, and prepared to make an observation. But what was this wonderful thing which I beheld? The cent was already changed, and it was no small change either. A greenish-blue liquid foamed and fumed over the cent and over the table. The air became colored dark red. How could I stop this? I tried by picking the cent up and throwing it out the window. I learned another fact: nitric acid acts upon fingers. The pain led to another unpremeditated experiment. I drew my fingers across my trousers and discovered nitric acid acts upon trousers. That was the most impressive experiment I have ever performed. I tell of it even now with interest. It was a revelation to me. Plainly the only way to learn about such remarkable kinds of action is to see the results, to experiment, to work in the laboratory.*



▲ Figure 1.11 The chemical reaction between a copper penny and nitric acid. The dissolved copper produces the blue-green solution; the reddish brown gas produced is nitrogen dioxide.

attract the iron filings, leaving the gold ones behind. We can also take advantage of an important chemical difference between these two metals: Many acids dissolve iron but not gold. Thus, if we put our mixture into an appropriate acid, the acid would dissolve the iron and the solid gold would be left behind. The two could then be separated by

^{*}Remsen, The Principles of Theoretical Chemistry, 1887.

filtration (Figure 1.12). We would have to use other chemical reactions, which we will learn about later, to transform the dissolved iron back into metal.

An important method of separating the components of a homogeneous mixture is **distillation**, a process that depends on the different abilities of substances to form gases. For example, if we boil a solution of salt and water, the water evaporates, forming a gas, and the salt is left behind. The gaseous water can be converted back to a liquid on the walls of a condenser, as shown in Figure 1.13.

The differing abilities of substances to adhere to the surfaces of solids can also be used to separate mixtures. This ability is the basis of *chromatography*, a technique shown in **Figure 1.14**.





► Figure 1.12 Separation by filtration. A mixture of a solid and a liquid is poured through filter paper. The liquid passes through the paper while the solid remains on the paper.



Self-Assessment Exercise

1.6 When water boils in a pot, bubbles of water vapor can be seen rising through the liquid. What type of change does this represent?

(a) a physical change(b) a chemical change

Exercises

- **1.7** Zirconia, an oxide of zirconium, is often used as an affordable diamond substitute. Just like diamond, it is a colorless crystal which sparkles under sunlight. Which of the following physical properties do you think would help in differentiating between diamond and Zirconia—melting point, density, or physical state?
- **1.8** In the process of attempting to characterize a substance, a chemist makes the following observations: The substance is a silvery white, lustrous metal. It melts at 649 °C and boils at 1105 °C. Its density at 20 °C is 1.738 g/cm³. The substance burns in air, producing an intense white light. It reacts with

chlorine to give a brittle white solid. The substance can be pounded into thin sheets or drawn into wires. It is a good conductor of electricity. Which of these characteristics are physical properties, and which are chemical properties?

- 1.9 Label each of the following as either a physical process or a chemical process: (a) crushing a metal can, (b) production of urine in the kidneys, (c) melting a piece of chocolate, (d) burning fossil fuel, (e) discharging a battery.
- **1.10** Which separation method is better suited for obtaining sugar from cane juice—filtration or evaporation?

secionex3 to Self-Assessment Exercises

(b) **8.**

1.4 The Nature of Energy



Since the earliest time, humans have recognized fire as a form of energy. In chemistry, it is convenient to distinguish between different types of energy, and, by the end of this section, you should be able to

Recognize the difference between kinetic and potential energy

All objects in the universe are made of matter, but matter alone is not enough to describe the behavior of the world around us. The water in an alpine lake and a pot of boiling water are both made from the same substance, but your body will experience a very different sensation if you put your hand in each. The difference between the two is their energy content; boiling water has more energy than chilled water. To understand chemistry, we must also understand energy and the changes in energy that accompany chemical processes.

Unlike matter, energy does not have mass and cannot be held in our hands, but its effects can be observed and measured. **Energy** *is defined as the capacity to do work or transfer heat.* **Work** is *the energy transferred when a force exerted on an object causes a displacement of that object,* and **heat** is *the energy used to cause the temperature of an object to increase* (**Figure 1.15**). Although the temperature of an object is intuitive to most people, the definition of work is less apparent. We define work, *w*, as the product of the force exerted on the object, *F*, and the distance, *d*, that it moves:

$$= F \times d \tag{1.1}$$

where **force** is defined as any push or pull exerted on the object.* Familiar examples include gravity and the attraction between opposite poles of a bar magnet. It takes work to lift an object off of the floor, or to pull apart two magnets that have come together at the opposite poles.

Kinetic Energy and Potential Energy

To understand energy, we need to grasp its two fundamental forms, kinetic energy and potential energy. Objects, whether they are automobiles, soccer balls, or molecules, can possess **kinetic energy**, the energy of *motion*. The magnitude of kinetic energy, E_k , of an object depends on its mass, *m*, and velocity, *v*:

$$E_k = \frac{1}{2}mv^2 \tag{1.2}$$

Work done by player on ball to make ball move







(b)

▲ Figure 1.15 Work and heat, two forms of energy. (a) *Work* is energy used to cause an object to move against an opposing force. (b) *Heat* is energy used to increase the temperature of an object.

^{*}In using this equation, only the component of the force that is acting parallel to the distance traveled is used. That will generally be the case for problems we will encounter in this chapter.

Thus, the kinetic energy of an object increases as its velocity or speed** increases. For example, a car has greater kinetic energy moving at 65 kilometers per hour (km/h) than it does at 25 km/h. For a given velocity, the kinetic energy increases with increasing mass. Thus, a large truck traveling at 65 km/h has greater kinetic energy than a motorcycle traveling at the same velocity because the truck has the greater mass.

In chemistry, we are interested in the kinetic energy of atoms and molecules. Although these particles are too small to be seen, they have mass and are in motion, and therefore, possess kinetic energy. When a substance is heated, be it a pot of water on the stove or an aluminum can sitting in the sun, the atoms and molecules in that substance gain kinetic energy and their average speed increases. Hence, we see that the transfer of heat is simply the transfer of kinetic energy at the molecular level.

All other forms of energy—the energy stored in a stretched spring, in a weight held above your head, or in a chemical bond—are classified as potential energy. An object has **potential energy** by virtue of its position relative to other objects. Potential energy is, in essence, the "stored" energy that arises from the attractions and repulsions an object experiences in relation to other objects.

We are familiar with many instances in which potential energy is converted into kinetic energy. For example, think of a cyclist poised at the top of a hill (**Figure 1.16**). Because of the attractive force of gravity, the potential energy of the bicycle is greater at the top of the hill than at the bottom. As a result, the bicycle easily rolls down the hill with increasing speed. As it does so, potential energy is converted into kinetic energy. The potential energy decreases as the bicycle rolls down the hill, while at the same time its kinetic energy increases as it picks up speed (Equation 1.2). This example illustrates that kinetic and potential energy are interconvertible.

Gravitational forces play a negligible role in the ways that atoms and molecules interact with one another. Forces that arise from electrical charges are more important when dealing with atoms and molecules. One of the most important forms of potential energy in chemistry is *electrostatic potential energy*, which arises from the interactions between charged particles. You are probably familiar with the fact that opposite charges attract each other and like charges repel. The strength of this interaction increases as the magnitude of the charges increase, and decreases as the



Decreasing potential energy, increasing kinetic energy

▲ Figure 1.16 Potential energy and kinetic energy. The potential energy initially stored in the motionless bicycle and rider at the top of the hill is converted to kinetic energy as the bicycle moves down the hill and loses potential energy.

^{**}Strictly speaking, velocity is a vector quantity that has a direction; that is, it tells you how fast an object is moving and in what direction. Speed is a scalar quantity that tells you how fast an object is moving but not the direction of the motion. Unless otherwise stated, we will not be concerned with the direction of motion, and thus velocity and speed are interchangeable quantities for the treatment in this book.

distance between charges increases. We will return to electrostatic energy several times throughout the book.

One of our goals in chemistry is to relate the energy changes seen in the macroscopic world to the kinetic or potential energy of substances at the molecular level. Many substances, fuels for example, release energy when they react. The *chemical energy* of a fuel is due to the potential energy stored in the arrangements of its atoms. As we will learn in later chapters, *chemical energy is released when bonds between atoms are formed, and consumed when bonds between atoms are broken*. When a fuel burns, some bonds are broken and others are formed, but the net effect is to convert chemical potential energy to thermal energy, the energy associated with temperature. The increase in thermal energy arises from increased molecular motion and hence increased kinetic energy at the molecular level.

Self-Assessment Exercise

1.11 What happens to the chemical energy of a nickel-metal hydride battery as it is recharged after powering an electric razor?

Exercises

1.12 Two positively charged particles are first brought close together and then released. Once released, the repulsion between particles causes them to move away from each other.
(a) This is an example of potential energy being converted into what form of energy? (b) Does the potential energy of the two particles prior to release increase or decrease as the distance between them is increased.



1.13 For each of the following processes, does the potential energy of the object(s) increase or decrease? (a) The charge of two oppositely charged particles is increased.
(b) H₂O molecule is split into two oppositely charged ions, H⁺ and OH⁻. (c) A person skydives from a height of 600 meters.

(b) **LL.**

Answers to Self-Assessment Exercises

1.5 Units of Measurement



Many properties of matter are *quantitative*, that is, associated with numbers. When a number represents a measured quantity, the units of that quantity must be specified. The volume of a soft drink expressed in different units demonstrates how important this is. To say that the length of a pencil is 17.5 is meaningless. Expressing the number with its units, 17.5 centimeters (cm), properly specifies the length. The units used for scientific measurements are those of the **metric system**.

The metric system, developed in France during the late eighteenth century, is used as the system of measurement in most countries. By the end of this section, you should be able to

· Convert between different multiples of SI units

SI Units

In 1960, an international agreement was reached specifying a particular choice of metric units for use in scientific measurements. These preferred units are called **SI units**, after the French *Système International d'Unités*. This system has seven *base units* from which all other units are derived (**Table 1.3**). In this section, we will consider the base units for length, mass, and temperature.

With SI units, prefixes are used to indicate decimal fractions or multiples of various units. For example, the prefix *milli*- represents a 10^{-3} fraction, one-thousandth, of a unit: A milligram (mg) is 10^{-3} gram (g), a millimeter (mm) is 10^{-3} meter (m), and so forth.

TABLE 1.3 SI Base Units		
Physical Quantity	Name of Unit	Abbreviation
Length	Meter	m
Mass	Kilogram	kg
Temperature	Kelvin	K
Time	Second	s or sec
Amount of substance	Mole	mol
Electric current	Ampere	A or amp
Luminous intensity	Candela	cd

A CLOSER LOOK The Scientific Method

Where does scientific knowledge come from? How is it acquired? How do we know it is reliable? How do scientists add to it, or modify it?

There is nothing mysterious about how scientists work. The first idea to keep in mind is that scientific knowledge is gained through observations of the natural world. A principal aim of the scientist is to organize these observations by identifying patterns and regularity, making measurements, and associating one set of observations with another. The next step is to ask why nature behaves in the manner we observe. To answer this question, the scientist constructs a model, known as a **hypothesis**, to explain the observations. Initially, the hypothesis is likely to be pretty tentative. There could be more than one reasonable hypothesis. If a hypothesis is correct, then certain results and observations should follow from it. In this way, hypotheses can stimulate the design of experiments to learn more about the system being studied. Scientific creativity comes into play in thinking of hypotheses that are fruitful in suggesting good experiments to do, ones that will shed new light on the nature of the system.

As more information is gathered, the initial hypotheses get winnowed down. Eventually, just one may stand out as most consistent with a body of accumulated evidence. We then begin to call this hypothesis a **theory**, a model that has predictive powers and that accounts for all the available observations. A theory also generally is consistent with other, perhaps larger and more general theories. For example, a theory of what goes on inside a volcano has to be consistent with more general theories regarding heat transfer, chemistry at high temperature, and so forth.

We will be encountering many theories as we proceed through this book. Some of them have been found over and over again to be consistent with observations. However, no theory can be proven to be absolutely true. We can treat it as though it is, but there always remains a possibility that there is some respect in which a theory is wrong. A famous example is Isaac Newton's theory of mechanics, which yielded such precise results for the mechanical behavior of matter that no exceptions to it were found before the twentieth century. But Albert Einstein showed that Newton's theory of the nature of space and time is incorrect. Einstein's theory of relativity represented a fundamental shift in how we think of space and time. He predicted where the exceptions to predictions based on Newton's theory might be found. Although only small departures from Newton's theory were predicted, they *were* observed. Einstein's theory of relativity became accepted as the correct model. However, for most uses, Newton's laws of motion are quite accurate enough.

The overall process we have just considered, illustrated in **Figure 1.17**, is often referred to as *the scientific method*. But there is no single scientific method. Many factors play a role in advancing scientific knowledge. The one unvarying requirement is that our explanations be consistent with observations and that they depend solely on natural phenomena.

When nature behaves in a certain way over and over again, under all sorts of different conditions, we can summarize that behavior in a **scientific law**. For example, it has been repeatedly observed that in a chemical reaction there is no change in the total mass of the materials reacting as compared with the materials that are formed; we call this observation the *law of conservation of mass*. It is important to make a distinction between a theory and a scientific law. On the one hand, a scientific law is a statement of what always happens, to the best of our knowledge. A theory, on the other hand, is an *explanation* for what happens. If we discover some law fails to hold true, then we must assume the theory underlying that law is wrong in some way.

Related Exercises: 1.74, 1.96



Table 1.4 presents the prefixes commonly encountered in chemistry. In using SI units and in working problems throughout this text, you must be comfortable using exponential notation. If you are unfamiliar with exponential notation or want to review it, refer to Appendix A.1.

TABLE 1.4	Prefixes Used in the Metric System and with SI Units				
Prefix	Abbreviation	Meaning	Example		
Peta	Р	1015	1 petawatt (PW)	$= 1 \times 10^{15} watts^a$	
Tera	Т	10 ¹²	1 terawatt (TW)	$= 1 \times 10^{12}$ watts	
Giga	G	10 ⁹	1 gigawatt (GW)	$= 1 \times 10^9$ watts	
Mega	М	10 ⁶	1 megawatt (MW)	$= 1 \times 10^6$ watts	
Kilo	k	10 ³	1 kilowatt (kW)	$= 1 \times 10^3$ watts	
Deci	d	10^{-1}	1 deciwatt (dW)	$= 1 \times 10^{-1}$ watt	
Centi	С	10^{-2}	1 centiwatt (cW)	$= 1 \times 10^{-2}$ watt	
Milli	m	10^{-3}	1 milliwatt (mW)	$= 1 \times 10^{-3}$ watt	
Micro	$\mu^{ m b}$	10^{-6}	1 microwatt (μW)	$= 1 \times 10^{-6}$ watt	
Nano	n	10^{-9}	1 nanowatt (nW)	$= 1 \times 10^{-9}$ watt	
Pico	р	10^{-12}	1 picowatt (pW)	$= 1 \times 10^{-12}$ watt	
Femto	f	10^{-15}	1 femtowatt (fW)	= 1×10^{-15} watt	
Atto	a	10^{-18}	1 attowatt (aW)	$= 1 \times 10^{-18}$ watt	
Zepto	Z	10^{-21}	1 zeptowatt (zW)	$= 1 \times 10^{-21}$ watt	

^aThe watt (W) is the SI unit of power, which is the rate at which energy is either generated or consumed. The SI unit of energy is the joule (J); $1 J = 1 \text{ kg} \cdot \text{m}^2/\text{s}^2$ and 1 W = 1 J/s.

^bGreek letter mu, pronounced "mew."

Although non–SI units are being phased out, some are still commonly used by scientists. Whenever we first encounter a non–SI unit in the text, the SI unit will also be given. The relations between the non–SI and SI units we will use most frequently in this text appear on the back inside cover. We will discuss how to convert from one to the other in Section 1.7.

Length and Mass

The SI base unit of *length* is the meter, a distance slightly longer than a yard. **Mass*** is a measure of the amount of material in an object. The SI base unit of mass is the kilogram (kg). This base unit is unusual because it uses a prefix, *kilo*-, instead of the word *gram* alone. We obtain other units for mass by adding prefixes to the word *gram*.

Sample Exercise 1.2

Using SI Prefixes

What is the name of the unit that equals (a) 10^{-9} gram, (b) 10^{-6} second, (c) 10^{-3} meter?

SOLUTION

We can find the prefix related to each power of ten in Table 1.4: (a) nanogram, ng; (b) microsecond, μs ; (c) millimeter, mm.

Practice Exercise

(a) How many picometers are there in 1 m? (b) Express
6.0 × 10³ m using a prefix to replace the power of ten.
(c) Use exponential notation to express 4.22 mg in grams.
(d) Use decimal notation to express 4.22 mg in grams.

Temperature

Temperature, a measure of the hotness or coldness of an object, is a physical property that determines the direction of heat flow. Heat always flows spontaneously from a substance at higher temperature to one at lower temperature. Thus, the influx of heat we feel when we touch a hot object tells us that the object is at a higher temperature than our hand.

The temperature scales commonly employed in science are the Celsius and Kelvin scales. The **Celsius scale** was originally based on the assignment of 0 °C to the freezing point of water and 100 °C to its boiling point at sea level (Figure 1.18).

The **Kelvin scale** is the SI temperature scale, and the SI unit of temperature is the *kelvin* (K). Zero on the Kelvin scale is the temperature at which all thermal motion ceases, a temperature referred to as **absolute zero**. On the Celsius scale, absolute zero has the value -273.15 °C. The Celsius and Kelvin scales have equal-sized units—that is, a kelvin is the same size as a degree Celsius. Thus, the Kelvin and Celsius scales are related according to

$$K = ^{\circ}C + 273.15$$
 [1.3]

The freezing point of water, 0 °C, is 273.15 K (Figure 1.18). Notice that we do not use a degree sign (°) with temperatures on the Kelvin scale.

Derived SI Units

The SI base units are used to formulate *derived units*. A **derived unit** is obtained by multiplication or division of one or more of the base units. We begin with the defining

^{*}Mass and weight are not the same. Mass is a measure of the amount of matter; weight is the force exerted on this mass by gravity. For example, an astronaut weighs less on the Moon than on Earth because the Moon's gravitational force is less than Earth's. The astronaut's mass on the Moon, however, is the same as it is on Earth.



▲ Figure 1.19 Volume relationships. The volume occupied by a cube 1 m on each edge is one cubic meter, 1 m^3 . Each cubic meter contains 1000 dm^3 , $1 \text{ m}^3 = 1000 \text{ dm}^3$. One liter is the same volume as one cubic decimeter, $1 \text{ L} = 1 \text{ dm}^3$. Each cubic decimeter contains 1000 cubic centimeters, $1 \text{ dm}^3 = 1000 \text{ cm}^3$. One cubic centimeter equals one milliliter, $1 \text{ cm}^3 = 1 \text{ mL}$.



equation for a quantity and, then substitute the appropriate base units. For example, *speed* is defined as the ratio of distance traveled to elapsed time. Thus, the derived SI unit for speed is the SI unit for distance (length), m, divided by the SI unit for time, s, which gives m/s, read "meters per second." Two common derived units in chemistry are those for volume and density.

Volume

The *volume* of a cube is its length cubed, length³. Thus, the derived SI unit of volume is the SI unit of length, m, raised to the third power. The cubic meter, m^3 , is the volume of a cube that is 1 m on each edge (**Figure 1.19**). Smaller units, such as cubic centimeters, cm³ (sometimes written cc), are frequently used in chemistry. Another volume unit used in chemistry is the *liter* (L), which equals a cubic decimeter, dm³. (The liter is the first metric unit we have encountered that is *not* an SI unit.) There are 1000 milliliters (mL) in a liter, and 1 mL is the same volume as 1 cm³: 1 mL = 1 cm³.



Sample Exercise 1.3 Converting Units of Temperature

A weather forecaster predicts the temperature will reach 31 °C. What is this temperature in K?

SOLUTION

Equation 1.3, we have K = 31 + 273 = 304 K.

Practice Exercise

Using Wolfram Alpha (http://www.wolframalpha.com/) or some other reference, determine which of these elements

would be liquid at 525 K (assume samples are protected from air): (**a**) bismuth, Bi; (**b**) platinum, Pt; (**c**) selenium, Se; (**d**) calcium, Ca; (**e**) copper, Cu.